

1. What is NARICOVER Lozenges and what it is used for?



NARICOVER Lozenges contains Naringin, a flavonoid present in citrus fruits and has antiviral and anti-inflammatory properties for use in throat medications. These are used for the effective management of Sore Throat & Dry Cough associated with Viral Pharyngitis & Laryngitis.

The action of chewing the lozenge allows the active ingredients to work in the area of discomfort and also helps lubricate and soothe the painful area. This helps relieve the soreness and discomfort of mouth and throat caused by viral infections.

2. What you need to know before you take NARICOVER Lozenges?



This product is recommended for adults, children (over 6 years old) and the elderly. Children under 6 years of age should not use this product.

Do not take this product if

- You are having diabetes mellitus (High Blood Sugar problem)
- You have an intolerance to certain sugars such as, Sucralose.
 These would be normally confirmed by your doctor.

Tell your doctor or pharmacist if you

- Are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- Suffer or are suffering from any other throat problems.
- It is not recommended to use this product if you are planning for pregnancy, are pregnant or are breast feeding. Ask your doctor or pharmacist for advice before taking any medicine.
- Important information about some of the ingredients in this
 product: This product contains Sucralose. If you have been told by
 your doctor that you have an intolerance to some sugars, contact
 your doctor before taking this product.

3. How to take NARICOVER Lozenges?



Remove one pillow pack of lozenge from the printed carton. Open the pillow pack and place the lozenge into the mouth, allowing it to dissolve slowly. Do not crush or chew the lozenge with force. It may damage the teeth or oral cavity.

Allow the lozenge to dissolve in the mouth slowly and repeat the same every 2-3 hours if required in 24 hours. It is not recommended for children under 6 years to use this product. Remember that young children can choke on lozenges. If symptoms persist for more than 3 days or if you notice any unusual symptoms, please consult your doctor immediately. Do not consume this product in excess, as exceeding the daily recommended dose may result in gastric disturbance (upset stomach). In case of gastric disturbance, please consult your doctor or pharmacist immediately. If you forget to take this product, do not take a double dose to make up for a forgotten lozenge. Continue to use this product, ask your doctor or pharmacist.

4. Possible side effects



Like all medicines, **NARICOVER** Lozenges can cause side effects, although not everybody gets them. Side effects are mild namely headache & flushing sensation.

If you experience any hypersensitivity to this product, i.e. itching, red spots on your body, stop taking this product and consult your doctor or pharmacist. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.